

# Lecture: Matrix Completion

<http://bicmr.pku.edu.cn/~wenzw/bigdata2020.html>

Acknowledgement: this slides is based on Prof. Jure Leskovec and Prof. Emmanuel Candes's lecture notes

# Recommendation systems

京东商城 北京

你好, 请登录 [免费注册](#) 我的订单 我的京东 京东会员 企业采购 客户服务 网站导航 手机京东

全部分类

笔记本电脑 台式机 办公设备 电脑办公 数码相机 打印机

自营旗舰店 [wacom](#) [关注我](#)

DTC-133 万与创意数位屏 [点击查看](#)

PTH-660/K9 影拓 Pro 进阶版 [点击查看](#)

DTK-1661 创意液晶数位屏 [点击查看](#)

全部商品 入门级 影拓 影拓 Pro 新帝 平板电脑 礼盒套装 Bamboo 原装配件

电脑、办公 > 外设产品 > 手写板 > Wacom > WacomCTL-6100WL...

Wacom CTL-6100WL/E 4096级压感 中号蓝牙连接intuos系列数位板绘图板

原价: ¥1199.00 降价通知

促销: 满600元减20元, 满2000元减50元

商品详情: 4.1-4.3英寸自适应板, 办公绘图、读图、线框、电路图及配件等指定商品满7元返10元 详情 >>

配送至: 北京海淀区八里庄街道... 可配货 支持 京尊达: 99元免基础运费(2kg以内)

品牌: 索尼, Wacom京东自营旗舰店 提供售后服务, 有货 (外地跨区发货), 暂无法直接发货, 18:00前下单, 预计9月4日07:00 (周二)送达

重量: 0.77kg

选择颜色: CTL6100 (5号) CTL6100WL (5号蓝牙版黄色) CTL6100WL (5号蓝牙版绿色) CTL6100 (4号) CTL6100WL (4号蓝牙版黄色) CTL6100WL (4号蓝牙版绿色) CTL6100WL (4号蓝牙版黄色) CTL6100WL (4号蓝牙版绿色)

增值服务: 2年免费换 ¥58 2年质保 ¥28 1年质保 ¥49

分期付款: 不分期 ¥399.66x3期 ¥205.85起x6期 ¥105.88起x12期 ¥55.92起x24期

加入购物车

加入购物车优惠

详细描述: 支持7光笔功能使用

店长推荐

¥509.00 ¥3899.00 ¥3199.00 ¥4499.00 ¥6349.00 ¥2499.00

Wacom京东自营旗舰店 [商品介绍](#) [规格与包装](#) [售后服务](#) [商品评价\(4.2万+\)](#) [商品问答](#) [加入购物车](#)

品牌: Wacom

加入购物车

商品名称: WacomCTL-6100WL/E-C 商品编号: 6973557 商品重量: 0.77kg 商品产地: 中国大陆

使用方式: 电磁笔 种类: 数位板 品牌: 影拓版 品牌: 409846

更多参数 >>

产品信息

产品特色

产品功能

## References:


<http://bicmr.pku.edu.cn/~wenzw/bigdata/07-recsys1.pdf>

<http://bicmr.pku.edu.cn/~wenzw/bigdata/08-recsys2.pdf>

# The Netflix Prize

- Training data
  - 100 million ratings, 480,000 users, 17,770 movies
  - 6 years of data: 2000-2005
- Test data
  - Last few ratings of each user (2.8 million)
  - Evaluation criterion: root mean squared error (RMSE):  
 $\sqrt{\sum_{xi} (r_{xi} - r_{xi}^*)^2}$ :  $r_{xi}$  and  $r_{xi}^*$  are the predicted and true rating of  $x$  on  $i$
  - Netflix Cinematch RMSE: 0.9514
- Competition
  - 2700+ teams
  - \$1 million prize for 10% improvement on Cinematch

# Netflix: evaluation



# Collaborative Filtering: weighted sum model

$$\hat{r}_{xi} = b_{xi} + \sum_{j \in N(i;x)} w_{ij}(r_{xj} - b_{xj})$$

- baseline estimate for  $r_{xi}$ :  $b_{xi} = \mu + b_x + b_i$   
 $\mu$ : overall mean rating  
 $b_x$ : rating deviation of user  $x$  = (avg. rating of user  $x$ ) -  $\mu$   
 $b_i$ : (avg. rating of movie  $i$ ) -  $\mu$
- We sum over all movies  $j$  that are similar to  $i$  and were rated by  $x$
- $w_{ij}$  is the interpolation weight (some real number). We allow:  
 $\sum_{j \in N(i,x)} w_{ij} \neq 1$
- $w_{ij}$  models interaction between pairs of movies (it does not depend on user  $x$ )
- $N(i;x)$ : set of movies rated by user  $x$  that are similar to movie  $i$

## Finding weights $w_{ij}$ ?

Find  $w_{ij}$  such that they work well on known (user, item) ratings:

$$\min_{w_{ij}} F(w) := \sum_x \left( \left[ b_{xi} + \sum_{j \in N(i;x)} w_{ij}(r_{xj} - b_{xj}) \right] - r_{xi} \right)^2$$

- Unconstrained optimization: quadratic function


$$\nabla_{w_{ij}} F(w) = 2 \sum_x \left( \left[ b_{xi} + \sum_{k \in N(i;x)} w_{ik}(r_{xk} - b_{xk}) \right] - r_{xi} \right) (r_{xj} - b_{xj}) = 0$$

for  $j \in \{N(i, x), \forall i, x\}$

- Equivalent to solving a system of linear equations?
- Steepest gradient descent method:  $w^{k+1} = w^k - \tau \nabla F(w)$
- Conjugate gradient method

# Latent factor models

- low rank factorization on Netflix data:  $R \approx Q \cdot P^T$



- For now let's assume we can approximate the rating matrix  $R$  as a product of "thin"  $Q \cdot P^T$   
 $R$  has missing entries but let's ignore that for now! Basically, we will want the reconstruction error to be small on known ratings and we don't care about the values on the missing ones

# Ratings as products of factors

- How to estimate the missing rating of user x for item i?

$$\hat{r}_{xi} = q_i \cdot p_x^T = \sum_f q_{if} p_{xf},$$

where  $q_i$  is row  $i$  of  $Q$  and  $p_x$  is column  $x$  of  $P^T$

	.1	-.4	.2
	<b>-.5</b>	<b>.6</b>	<b>.5</b>
	-.2	.3	.5
	1.1	2.1	.3
	-.7	2.1	-.2
	-.1	.7	.3
items			
	f factors		

	users											
	1.1	-.2	.3	.5	<b>-.2</b>	-.5	.8	-.4	.3	1.4	2.4	-.9
	-.8	.7	.5	1.4	<b>.3</b>	-.1	1.4	2.9	-.7	1.2	-.1	1.3
	2.1	-.4	.6	1.7	<b>2.4</b>	.9	-.3	.4	.8	.7	-.6	.1
f factors												
	$P^T$											



# Latent factor models

- Minimize SSE on training data!
- Use specialized methods to find P, Q such that  $\hat{r}_{xi} = q_i \cdot p_x^T$

$$\min_{P, Q} \sum_{(i,x) \in \text{training}} (r_{xi} - q_i \cdot p_x^T)^2$$

We don't require cols of P, Q to be orthogonal/unit length

- P, Q map users/movies to a latent space
- Add regularization:

$$\min_{P, Q} \sum_{(i,x) \in \text{training}} (r_{xi} - q_i \cdot p_x^T)^2 + \lambda \left[ \sum_x \|p_x\|_2^2 + \sum_i \|q_i\|_2^2 \right]$$

$\lambda$  is called regularization parameters

# Gradient descent method

$$\min_{P,Q} F(P, Q) := \sum_{(i,x) \in \text{training}} (r_{xi} - q_i \cdot p_x^T)^2 + \lambda \left[ \sum_x \|p_x\|_2^2 + \sum_i \|q_i\|_2^2 \right]$$

Gradient decent:

- Initialize P and Q (using SVD, pretend missing ratings are 0)
- Do gradient descent:  
 $P^{k+1} \leftarrow P^k - \tau \nabla_P F(P^k, Q^k),$   
 $Q^{k+1} \leftarrow Q^k - \tau \nabla_Q F(P^k, Q^k),$   
where  $(\nabla_Q F)_{if} = -2 \sum_{x,i} (r_{xi} - q_i p_x^T) p_{xf} + 2\lambda q_{if}$ . Here  $q_{if}$  is entry f of row  $q_i$  of matrix Q
- Computing gradients is slow when the dimension is huge

# Stochastic gradient descent method

Observation: Let  $q_{if}$  be entry  $f$  of row  $q_i$  of matrix  $Q$

$$(\nabla_Q F)_{if} = \sum_{x,i} (-2(r_{xi} - q_i p_x^T) p_{xf} + 2\lambda q_{if}) = \sum_{x,i} \nabla_Q F(r_{xi})$$

$$(\nabla_P F)_{xf} = \sum_{x,i} (-2(r_{xi} - q_i p_x^T) q_{xf} + 2\lambda p_{if}) = \sum_{x,i} \nabla_P F(r_{xi})$$

Stochastic gradient decent:

- Instead of evaluating gradient over all ratings, evaluate it for each individual rating and make a step
- $P \leftarrow P - \tau \nabla_P F(r_{xi})$   
 $Q \leftarrow Q - \tau \nabla_Q F(r_{xi})$
- Need more steps but each step is computed much faster

# Latent factor models with biases

predicted models:

$$\hat{r}_{xi} = \mu + b_x + b_i + q_i \cdot p_x^T$$

$\mu$ : overall mean rating,  $b_x$ : Bias for user  $x$ ,  $b_i$ : Bias for movie  $i$


New model:

$$\min_{P, Q, b_x, b_i} \sum_{(i,x) \in \text{training}} (r_{xi} - (\mu + b_x + b_i + q_i \cdot p_x^T))^2 + \lambda \left[ \sum_x \|p_x\|_2^2 + \sum_i \|q_i\|_2^2 + \|b_x\|_2^2 + \|b_i\|_2^2 \right]$$

- Both biases  $b_x$ ,  $b_i$  as well as interactions  $q_i$ ,  $p_x$  are treated as parameters (we estimate them)
- Add time dependence to biases:

$$\hat{r}_{xi} = \mu + b_x(t) + b_i(t) + q_i \cdot p_x^T$$

# Netflix: performance



# Netflix: performance

## Netflix Prize

COMPLETED

[Home](#) [Rules](#) [Leaderboard](#) [Update](#) [Download](#)

### Leaderboard

Showing Test Score. [Click here to show quiz score](#)

Display top  leaders.

Rank	Team Name	Best Test Score	% Improvement	Best Submit Time
------	-----------	-----------------	---------------	------------------

Grand Prize - RMSE = 0.8567 - Winning Team: BellKor's Pragmatic Chaos

1	<a href="#">BellKor's Pragmatic Chaos</a>	0.8567	10.06	2009-07-26 18:18:28
2	<a href="#">The Ensemble</a>	0.8567	10.06	2009-07-26 18:38:22
3	<a href="#">Grand Prize Team</a>	0.8562	0.95	2009-07-10 21:24:40
4	<a href="#">Opera Solutions and Vandelay United</a>	0.8588	9.84	2009-07-10 01:12:31
5	<a href="#">Vandelay Industries I</a>	0.8591	9.81	2009-07-10 00:32:20
6	<a href="#">PragmaticTheory</a>	0.8594	9.77	2009-06-24 12:06:56
7	<a href="#">BellKor in BigChaos</a>	0.8601	9.70	2009-05-13 08:14:09
8	<a href="#">Dace</a>	0.8612	9.59	2009-07-24 17:18:43
9	<a href="#">Feeds2</a>	0.8622	9.48	2009-07-12 13:11:51
10	<a href="#">BigChaos</a>	0.8623	9.47	2009-04-07 12:33:59
11	<a href="#">Opera Solutions</a>	0.8623	9.47	2009-07-24 00:34:07
12	<a href="#">BellKor</a>	0.8624	9.46	2009-07-26 17:19:11

Progress Prize 2008 - RMSE = 0.8627 - Winning Team: BellKor in BigChaos

13	<a href="#">vionellone</a>	0.8642	0.27	2009-07-15 14:53:22
----	----------------------------	--------	------	---------------------

# General matrix completion

# Matrix completion

- Matrix  $M \in \mathbb{R}^{n_1 \times n_2}$
- Observe subset of entries
- Can we guess the missing entries?

$$\begin{bmatrix} \times & ? & ? & ? & \times & ? \\ ? & ? & \times & \times & ? & ? \\ \times & ? & ? & \times & ? & ? \\ ? & ? & \times & ? & ? & \times \\ \times & ? & ? & ? & ? & ? \\ ? & ? & \times & \times & ? & ? \end{bmatrix}$$



# Which algorithm ?

Hope: only **one** low-rank matrix consistent with the sampled entries

Recovery by minimum complexity

$$\begin{array}{ll} \text{minimize} & \text{rank}(X) \\ \text{subject to} & X_{ij} = M_{ij}, \quad (i,j) \in \Omega \end{array}$$

## Problem

- This is NP-hard
- Doubly exponential in  $n$  (?)

# SVD - Properties

## Theorem: SVD

If  $A$  is a real  $m$ -by- $n$  matrix, then there exists

$$U = [u_1, \dots, u_m] \in \mathbb{R}^{m \times m} \text{ and } V = [v_1, \dots, v_n] \in \mathbb{R}^{n \times n}$$

such that  $U^T U = I$ ,  $V^T V = I$  and

$$U^T A V = \text{diag}(\sigma_1, \dots, \sigma_p) \in \mathbb{R}^{m \times n}, \quad p = \min(m, n),$$

where  $\sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_p \geq 0$ .

- Proof: Let  $V_1 \in \mathbb{R}^{n \times r}$  has orthonormal columns, then exists  $V_2 \in \mathbb{R}^{n \times (n-r)}$  such that  $V = [V_1, V_2]$  is orthogonal.
- Let  $x \in \mathbb{R}^n$  and  $y \in \mathbb{R}^m$  be unit 2-norm vectors:  $Ax = \sigma y$  with  $\sigma = \|A\|_2$ . Then exists  $V_2 \in \mathbb{R}^{n \times (n-1)}$  and  $U_2 \in \mathbb{R}^{m \times (m-1)}$  so  $V = [x, V_2] \in \mathbb{R}^{n \times n}$  and  $U = [y, U_2] \in \mathbb{R}^{m \times m}$  are orthogonal.

- Then it can be proved that  $U^T AV$  has the following structure

$$U^T AV = \begin{pmatrix} \sigma & w^T \\ 0 & B \end{pmatrix} \equiv A_1.$$

Since

$$\left\| A_1 \begin{pmatrix} \sigma \\ w \end{pmatrix} \right\|_2^2 \geq (\sigma^2 + w^T w)^2,$$

we have  $\|A_1\|_2^2 \geq (\sigma^2 + w^T w)$ . But  $\sigma^2 = \|A\|_2^2 = \|A_1\|_2^2$ , and so we must have  $w = 0$ . An induction gives the proof.

Properties:

- $AV = U\Sigma$ ,  $A^T U = V\Sigma^T$ :  $Av_i = \sigma u_i$ ,  $A^T u_i = \sigma_i v_i$ ,  $i = 1, \dots, p$ .
- $\text{rank}(A) = r$ ,  $\text{null}(A) = \text{span}\{v_{r+1}, \dots, v_n\}$ ,  $\text{ran}(A) = \text{span}\{u_1, \dots, u_r\}$
- $A = \sum_{i=1}^r \sigma_i u_i v_i^T$
- $\|A\|_F^2 = \sigma_1^2 + \dots + \sigma_p^2$ ,  $\|A\|_2 = \sigma_1$

# SVD - Best Low Rank Approximation

## Theorem

Let the SVD of  $A \in \mathbb{R}^{m \times n}$  be given in Theorem: SVD. If  $k < r = \text{rank}(A)$  and  $A_k = \sum_{i=1}^k \sigma_i u_i v_i^T$ , then

$$\min_{\text{rank}(B)=k} \|A - B\|_2 = \|A - A_k\|_2 = \sigma_{k+1}.$$

- Proof: Since  $U^T A_k V = \text{diag}(\sigma_1, \dots, \sigma_k, 0, \dots, 0)$  it follows that  $\text{rank}(A_k) = k$  and  $U^T (A - A_k) V = \text{diag}(0, \dots, 0, \sigma_{k+1}, \dots, \sigma_p)$ . Hence  $\|A - A_k\|_2 = \sigma_{k+1}$ .
- Suppose  $\text{rank}(B) = k$  for some  $B \in \mathbb{R}^{m \times n}$ . We can find orthonormal vectors  $x_1, \dots, x_{n-k}$  so  $\text{null}(B) = \text{span}\{x_1, \dots, x_{n-k}\}$ . A dimension argument shows:

$$\text{span}\{x_1, \dots, x_{n-k}\} \cap \text{span}\{v_1, \dots, v_{k+1}\} \neq \{0\}$$

- Let  $z$  be a unit 2-norm vector in this intersection. Since  $Bz = 0$  and

$$Az = \sum_{i=1}^{k+1} \sigma_i (v_i^T z) u_i,$$

we have

$$\|A - B\|_2^2 \geq \|(A - B)z\|_2^2 = \|Az\|_2^2 = \sum_{i=1}^{k+1} \sigma_i^2 (v_i^T z)^2 \geq \sigma_{k+1}^2$$

Comments:

- So zeroing small  $\sigma_i$  introduces less error
- How many  $\sigma$ s to keep? Rule of thumb: keep 80-90% of 'energy' ( $= \sum \sigma_i^2$ )

# SVD - Complexity

- To compute SVD:  $O(nm^2)$  or  $O(n^2m)$
- But:
  - Less work, if we just want singular values
  - or if we want first  $k$  singular vectors
  - or if the matrix is sparse
- Implemented in linear algebra packages like
  - Dense matrix: LAPACK
  - Sparse Matrix: ARPACK, PROPACK
  - High Level Software packages: Matlab, SPlus, Mathematica ...

# Relation to Eigen-decomposition

- SVD gives us  $A = U\Sigma V^\top$
- Eigen-decomposition:  $A = X\Lambda X^\top$ 
  - $A$  is symmetric
  - $U, V, X$  are orthonormal
  - $\Lambda, \Sigma$  are diagonal
- $AA^\top = U\Sigma\Sigma^\top U^\top$
- $A^\top A = V\Sigma\Sigma^\top V^\top$
- $\lambda_i(A^\top A) = \sigma_i^2(A)$

# Nuclear-norm minimization

Singular value decomposition

$$X = \sum_{k=1}^r \sigma_k u_k v_k^*$$

- $\{\sigma_k\}$ : singular values,  $\{u_k\}, \{v_k\}$ : singular vectors

Nuclear norm ( $\sigma_i(X)$  is  $i$ th largest singular value of  $X$ )

$$\|X\|_* = \sum_{i=1}^n \sigma_i(X)$$

Heuristic

$$\begin{array}{ll} \text{minimize} & \|X\|_* \\ \text{subject to} & X_{ij} = M_{ij}, \quad (i,j) \in \Omega \end{array}$$

- Convex relaxation of the rank minimization program



# Connections with compressed sensing

## General setup

Rank minimization

$$\begin{array}{ll}\text{minimize} & \text{rank}(X) \\ \text{subject to} & \mathcal{A}(X) = b\end{array}$$

Convex relaxation

$$\begin{array}{ll}\text{minimize} & \|X\|_* \\ \text{subject to} & \mathcal{A}(X) = b\end{array}$$

Suppose  $X = \text{diag}(x), x \in \mathbb{R}^n$

- $\text{rank}(X) = \sum_i 1_{(x_i \neq 0)} = \|x\|_{\ell_0}$
- $\|X\|_* = \sum_i |x_i| = \|x\|_{\ell_1}$

Rank minimization

$$\begin{array}{ll}\text{minimize} & \|x\|_{\ell_0} \\ \text{subject to} & Ax = b\end{array}$$

Convex relaxation

$$\begin{array}{ll}\text{minimize} & \|x\|_{\ell_1} \\ \text{subject to} & Ax = b\end{array}$$

This is compressed sensing!

# Correspondence

parsimony concept	cardinality	rank
Hilbert Space norm	Euclidean	Frobenius
sparsity inducing norm	$\ell_1$	nuclear
dual norm	$\ell_\infty$	operator
norm additivity	disjoint support	orthogonal row and column spaces
convex optimization	linear programming	semidefinite programming

Table: From Recht Parrilo Fazel (08)

# Semidefinite programming (SDP)

- Special class of convex optimization problems
- Relatively natural extension of linear programming (LP)
- “Efficient” numerical solvers (interior point methods)

LP:  $x \in \mathbb{R}^n$

minimize  $\langle c, x \rangle$   
subject to  $Ax = b$   
 $x \geq 0$

SDP:  $X \in \mathbb{R}^{n \times n}$

minimize  $\langle C, X \rangle$   
subject to  $\langle A_k, X \rangle = b_k$   
 $X \succcurlyeq 0$

Standard inner product:  $\langle C, X \rangle = \text{trace}(C^*X)$

# SOCP/SDP Duality

$$\begin{array}{ll} \text{(P)} & \min \quad c^\top x \\ & \text{s.t.} \quad Ax = b, x_{\mathcal{Q}} \succeq 0 \end{array}$$

$$\begin{array}{ll} \text{(D)} & \max \quad b^\top y \\ & \text{s.t.} \quad A^\top y + s = c, s_{\mathcal{Q}} \succeq 0 \end{array}$$

$$\begin{array}{ll} \text{(P)} & \min \quad \langle C, X \rangle \\ & \text{s.t.} \quad \langle A_1, X \rangle = b_1 \\ & \quad \dots \\ & \quad \langle A_m, X \rangle = b_m \\ & \quad X \succeq 0 \end{array}$$

$$\begin{array}{ll} \text{(D)} & \max \quad b^\top y \\ & \text{s.t.} \quad \sum_i y_i A_i + S = C \\ & \quad S \succeq 0 \end{array}$$

## Strong duality

- If  $p^* > -\infty$ , (P) is **strictly** feasible, then (D) is feasible and  $p^* = d^*$
- If  $d^* < +\infty$ , (D) is **strictly** feasible, then (P) is feasible and  $p^* = d^*$
- If (P) and (D) has **strictly** feasible solutions, then both have optimal solutions.

# Semidefinite program

$$(D) \quad \begin{array}{ll} \min & -b^\top y \\ \text{s.t.} & y_1 A_1 + \dots + y_m A_m \preceq C \end{array}$$

- $A_i, C \in \mathcal{S}^k$ , multiplier is matrix  $X \in \mathcal{S}^k$
- Lagrangian  $\mathcal{L}(y, X) = -b^\top y + \langle X, y_1 A_1 + \dots + y_m A_m - C \rangle$
- dual function

$$g(X) = \inf_y \mathcal{L}(y, X) = \begin{cases} -\langle C, X \rangle, & \langle A_i, X \rangle = b_i \\ -\infty & \text{otherwise} \end{cases}$$

The dual of (D) is

$$\begin{array}{ll} \min & \langle C, X \rangle \\ \text{s.t.} & \langle A_i, X \rangle = b_i, X \succeq 0 \end{array}$$

$p^* = d^*$  if primal SDP is strictly feasible.

# SDP Relaxation of Maxcut

$$\begin{array}{lll} \min & x^\top W x & \\ \text{s.t.} & x_i^2 = 1 & \end{array} \quad \Rightarrow \quad \begin{array}{ll} \max & -1^\top \nu \\ \text{s.t.} & W + \text{diag}(\nu) \succeq 0 \end{array} \quad \Longleftrightarrow \quad \begin{array}{ll} \min & \text{Tr}(WX) \\ \text{s.t.} & X_{ii} = 1 \\ & X \succeq 0 \end{array}$$

- a nonconvex problem; feasible set contains  $2^n$  discrete points
- interpretation: partition  $\{1, \dots, n\}$  in two sets;  $W_{ij}$  is cost of assigning  $i, j$  to the same set;  $-W_{ij}$  is cost of assigning to different sets

dual function

$$\begin{aligned} g(\nu) &= \inf_x \left( x^\top W x + \sum_i \nu_i (x_i^2 - 1) \right) = \inf_x x^\top (W + \text{diag}(\nu)) x - 1^\top \nu \\ &= \begin{cases} -1^\top \nu & W + \text{diag}(\nu) \succeq 0 \\ -\infty & \text{otherwise} \end{cases} \end{aligned}$$

# Positive semidefinite unknown: SDP formulation

Suppose unknown matrix  $X$  is positive semidefinite

$$\begin{array}{ll} \min & \sum_{i=1}^n \sigma_i(X) \\ \text{s.t.} & X_{ij} = M_{ij} \quad (i,j) \in \Omega \\ & X \succcurlyeq 0 \end{array} \quad \Leftrightarrow \quad \begin{array}{ll} \min & \text{trace}(X) \\ \text{s.t.} & X_{ij} = M_{ij} \quad (i,j) \in \Omega \\ & X \succcurlyeq 0 \end{array}$$

Trace heuristic: Mesbahi & Papavassilopoulos (1997), Beck & D'Andrea (1998)

# General SDP formulation

Let  $X \in \mathbb{R}^{m \times n}$ . For a given norm  $\|\cdot\|$ , the dual norm  $\|\cdot\|_d$  is defined as

$$\|X\|_d := \sup\{\langle X, Y \rangle : Y \in \mathbb{R}^{m \times n}, \|Y\| \leq 1\}$$

Nuclear norm and spectral norms are dual:

$$\|X\| := \sigma_1(X), \quad \|X\|_* = \sum_i \sigma_i(X).$$

$$\begin{aligned} \text{(P)} \quad & \max_Y \langle X, Y \rangle \\ & \text{s.t. } \|Y\|_2 \leq 1 \end{aligned} \quad \Leftrightarrow \quad \begin{aligned} & \max_Y 2\langle X, Y \rangle \\ & \text{s.t. } \begin{bmatrix} I_m & Y \\ Y^\top & I_n \end{bmatrix} \succcurlyeq 0 \end{aligned} \quad \Leftrightarrow \quad \begin{aligned} & \min_Z - \left\langle Z, \begin{bmatrix} 0 & X \\ X^\top & 0 \end{bmatrix} \right\rangle \\ & \text{s.t. } Z_1 = I_m \\ & \quad Z_2 = I_n \\ & \quad Z = \begin{bmatrix} Z_1 & Z_3 \\ Z_3^\top & Z_2 \end{bmatrix} \succcurlyeq 0 \end{aligned}$$



# General SDP formulation

The Lagrangian dual problem is:

$$\max_{W_1, W_2} \min_{Z \succeq 0} - \left\langle Z, \begin{bmatrix} 0 & X \\ X^* & 0 \end{bmatrix} \right\rangle + \langle Z_1 - I_m, W_1 \rangle + \langle Z_2 - I_n, W_2 \rangle$$

*strong duality* after a scaling of 1/2 and change of variables  $X$  to  $-X$

$$\begin{aligned} \text{(D)} \quad & \text{minimize} \quad \frac{1}{2} (\text{trace}(W_1) + \text{trace}(W_2)) \\ & \text{subject to} \quad \begin{bmatrix} W_1 & X \\ X^\top & W_2 \end{bmatrix} \succeq 0 \end{aligned}$$

Optimization variables:  $W_1 \in \mathbb{R}^{n_1 \times n_1}$ ,  $W_2 \in \mathbb{R}^{n_2 \times n_2}$ .

Proposition 2.1 in "Guaranteed Minimum-Rank Solutions of Linear Matrix Equations via Nuclear Norm Minimization", Benjamin Recht, Maryam Fazel, Pablo A. Parrilo

# General SDP formulation

## Nuclear norm minimization

$$\begin{array}{ll} \min \|X\|_* \\ \text{s.t. } \mathcal{A}(X) = b \end{array} \iff \begin{array}{ll} \max b^\top y \\ \text{s.t. } \|\mathcal{A}^*(y)\| \leq 1 \end{array}$$

## SDP Reformulation

$$\begin{array}{ll} \min \frac{1}{2} (\text{trace}(W_1) + \text{trace}(W_2)) \\ \text{s.t. } \mathcal{A}(X) = b \\ \begin{bmatrix} W_1 & X \\ X^\top & W_2 \end{bmatrix} \succcurlyeq 0 \end{array} \iff \begin{array}{ll} \max b^\top y \\ \text{s.t. } \begin{bmatrix} I & \mathcal{A}^*(y) \\ (\mathcal{A}^*(y))^\top & I \end{bmatrix} \succcurlyeq 0 \end{array}$$

# Matrix recovery

$$M = \sum_{k=1}^2 \sigma_k u_k u_k^*, \quad \begin{aligned} u_1 &= (e_1 + e_2)/\sqrt{2}, \\ u_2 &= (e_1 - e_2)/\sqrt{2} \end{aligned}$$

$$M = \begin{bmatrix} * & * & 0 & \dots & 0 & 0 \\ * & * & 0 & \dots & 0 & 0 \\ 0 & 0 & 0 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & 0 & 0 \end{bmatrix}$$

Cannot be recovered from a small set of entries

Rank-1 matrix  $M = xy^*$

$$M_{ij} = x_i y_j^*$$

$$\begin{bmatrix} \times & \times & \times & \times & \times & \times \\ \times & \times & \times & \times & \times & \times \\ \times & \times & \times & \times & \times & \times \\ \times & \times & \times & \times & \times & \times \\ \times & \times & \times & \times & \times & \times \end{bmatrix}$$

If single row (or column) is not sampled  $\rightarrow$  recovery is not possible

*What happens for almost all sampling sets?*

$\Omega$  subset of  $m$  entries selected uniformly at random

# References

- Jianfeng Cai, Emmanuel Candes, Zuowei Shen, *Singular value thresholding algorithm for matrix completion*
- Shiqian Ma, Donald Goldfarb, Lifeng Chen, *Fixed point and Bregman iterative methods for matrix rank minimization*
- Zaiwen Wen, Wotao Yin, Yin Zhang, *Solving a low-rank factorization model for matrix completion by a nonlinear successive over-relaxation algorithm*
- Onureena Banerjee, Laurent El Ghaoui, Alexandre d'Aspremont, *Model Selection Through Sparse Maximum Likelihood Estimation for Multivariate Gaussian or Binary Data*
- Zhaosong Lu, *Smooth optimization approach for sparse covariance selection*

# Matrix Rank Minimization

Given  $X \in \mathbb{R}^{m \times n}$ ,  $\mathcal{A} : \mathbb{R}^{m \times n} \rightarrow \mathbb{R}^p$ ,  $b \in \mathbb{R}^p$ , we consider

- the matrix rank minimization problem:

$$\min \text{rank}(X), \text{ s.t. } \mathcal{A}(X) = b$$

- matrix completion problem:

$$\min \text{rank}(X), \text{ s.t. } X_{ij} = M_{ij}, (i, j) \in \Omega$$

- nuclear norm minimization:

$$\min \|X\|_* \text{ s.t. } \mathcal{A}(X) = b$$

where  $\|X\|_* = \sum_i \sigma_i$  and  $\sigma_i = i$ th singular value of matrix  $X$ .

# Quadratic penalty framework

- Unconstrained Nuclear Norm Minimization:

$$\min F(X) := \mu \|X\|_* + \frac{1}{2} \|\mathcal{A}(X) - b\|_2^2.$$

- Optimality condition:

$$\mathbf{0} \in \mu \partial \|X^*\|_* + \mathcal{A}^*(\mathcal{A}(X^*) - b),$$

where  $\partial \|X\|_* = \{UV^\top + W : U^\top W = 0, WV = 0, \|W\|_2 \leq 1\}$ .

- Linearization approach ( $g$  is the gradient of  $\frac{1}{2} \|\mathcal{A}(X) - b\|_2^2$ ):

$$\begin{aligned} X^{k+1} &:= \arg \min_X \mu \|X\|_* + \langle g^k, X - X^k \rangle + \frac{1}{2\tau} \|X - X^k\|_F^2 \\ &= \arg \min_X \mu \|X\|_* + \frac{1}{2\tau} \|X - (X^k - \tau g^k)\|_F^2 \end{aligned}$$

# Matrix Shrinkage Operator

For a matrix  $Y \in \mathbb{R}^{m \times n}$ , consider:

$$\min_{X \in \mathbb{R}^{m \times n}} \nu \|X\|_* + \frac{1}{2} \|X - Y\|_F^2.$$

The optimal solution is:

$$X := S(Y, \nu) = U \text{Diag}(s(\sigma, \nu)) V^\top,$$

- SVD:  $Y = U \text{Diag}(\sigma) V^\top$
- Thresholding operator:

$$s(x, \nu) := \bar{x}, \text{ with } \bar{x}_i = \begin{cases} x_i - \nu, & \text{if } x_i - \nu > 0 \\ 0, & \text{o.w.} \end{cases}$$



# Fixed Point Method (Proximal gradient method)

## Fixed Point Iterative Scheme

$$\begin{cases} Y^k = X^k - \tau \mathcal{A}^*(\mathcal{A}(X^k) - b) \\ X^{k+1} = S(Y^k, \tau \mu). \end{cases}$$

**Lemma:** Matrix shrinkage operator is non-expansive. i.e.,

$$\|S(Y_1, \nu) - S(Y_2, \nu)\|_F \leq \|Y_1 - Y_2\|_F.$$

Complexity of the fixed point method:

$$F(X^k) - F(X^*) \leq \frac{L_f \|X^0 - X^*\|^2}{2k}$$

# Accelerated proximal gradient (APG) method

APG algorithm ( $t^{-1} = t^0 = 1$ ):

$$\begin{aligned}Y^k &= X^k + \frac{t^{k-1} - 1}{t^k}(X^k - X^{k-1}) \\G^k &= Y^k - (\tau^k)^{-1} \mathcal{A}^*(\mathcal{A}(Y^k) - b) \\X^{k+1} &= S_{\tau^k}(G^k), \quad t^{k+1} = \frac{1 + \sqrt{1 + 4(t^k)^2}}{2}\end{aligned}$$

Complexity:

$$F(X^k) - F(X^*) \leq \frac{2L_f \|X^0 - X^*\|^2}{(k+1)^2}$$

Linearized Bregman method:

$$\begin{aligned}V^{k+1} &:= V^k - \tau \mathcal{A}^*(\mathcal{A}(X^k) - b) \\X^{k+1} &:= S_{\tau\mu}(V^{k+1})\end{aligned}$$

Convergence to

$$\min \tau \|X\|_* + \frac{1}{2} \|X\|_F^2, \text{ s.t. } \mathcal{A}(X) = b$$

# Review of Bregman method

Consider the problem:

$$\min \|X\|_*, \quad \text{s.t. } \mathcal{A}(X) = b$$

Bregman method:

- $D^{P^k}(X, X^k) := \|X\|_* - \|X^k\|_* - \langle P^k, X - X^k \rangle$
- $X^{k+1} := \arg \min_X \mu D^{P^k}(X, X^k) + \frac{1}{2} \|\mathcal{A}(X) - b\|_2^2$
- $P^{k+1} = P^k + \frac{1}{\mu} \mathcal{A}^\top(b - \mathcal{A}(X^{k+1}))$

Augmented Lagrangian (updating multiplier or  $b$ ):

- $X^{k+1} := \arg \min_X \mu \|X\|_* + \frac{1}{2} \|\mathcal{A}(X) - b^k\|_2^2$
- $b^{k+1} = b + (b^k - \mathcal{A}(X^{k+1}))$

They are equivalent, see Yin-Osher-Goldfarb-Darbon

# Linearized approaches

## Linearized Bregman method:

$$\begin{aligned}X^{k+1} &:= \arg \min \mu D^{p^k}(X, X^k) + \left\langle \mathcal{A}^\top(\mathcal{A}(X^k) - b), X - X^k \right\rangle + \frac{1}{2\delta} \|X - X^k\|_F^2, \\P^{k+1} &:= P^k - \frac{1}{\mu\delta}(X^{k+1} - X^k) - \frac{1}{\mu}\mathcal{A}^\top(\mathcal{A}(X^k) - b),\end{aligned}$$

which is equivalent to

$$\begin{aligned}X^{k+1} &:= \arg \min \mu \|X\|_* + \frac{1}{2\delta} \|X - V^k\|_F^2 \\V^{k+1} &:= V^k - \delta \mathcal{A}^\top(\mathcal{A}(X^{k+1}) - b)\end{aligned}$$

## Bregmanized operator splitting:

$$\begin{aligned}X^{k+1} &:= \arg \min \mu \|X\|_* + \left\langle \mathcal{A}^\top(\mathcal{A}(X^k) - b^k), X - X^k \right\rangle + \frac{1}{2\delta} \|X - X^k\|_F^2 \\b^{k+1} &= b + (b^k - \mathcal{A}(X^{k+1}))\end{aligned}$$

Are they equivalent?

# Linearized approaches

## Linearized Bregman method:

$$\begin{aligned} X^{k+1} &:= \arg \min \mu D^{p^k}(X, X^k) + \left\langle \mathcal{A}^\top(\mathcal{A}(X^k) - b), X - X^k \right\rangle + \frac{1}{2\delta} \|X - X^k\|_F^2, \\ P^{k+1} &:= P^k - \frac{1}{\mu\delta}(X^{k+1} - X^k) - \frac{1}{\mu} \mathcal{A}^\top(\mathcal{A}(X^k) - b), \end{aligned}$$

which is equivalent to

$$\begin{aligned} X^{k+1} &:= \mathcal{S}(V^k, \mu\delta) \\ V^{k+1} &:= V^k - \delta \mathcal{A}^\top(\mathcal{A}(X^{k+1}) - b) \end{aligned} \quad \text{or} \quad \begin{aligned} X^{k+1} &:= \mathcal{S}(\delta \mathcal{A}^\top(b^k), \mu\delta) \\ b^{k+1} &:= b + (b^k - \mathcal{A}(X^{k+1})) \end{aligned}$$

## Bregmanized operator splitting:

$$\begin{aligned} X^{k+1} &:= \mathcal{S}(X^k - \delta(\mathcal{A}^\top(\mathcal{A}(X^k) - b^k)), \mu\delta) = \mathcal{S}(\delta \mathcal{A}^\top(b^k) + \textcolor{blue}{X}^k - \delta \mathcal{A}^\top(\mathcal{A}(X^k)), \mu\delta) \\ b^{k+1} &= b + (b^k - \mathcal{A}(X^{k+1})) \end{aligned}$$

# Low-rank factorization model

- Finding a low-rank matrix  $W$  so that  $\|\mathcal{P}_\Omega(W - M)\|_F^2$  or the distance between  $W$  and  $\{Z \in \mathbb{R}^{m \times n}, Z_{ij} = M_{ij}, \forall (i,j) \in \Omega\}$  is minimized.
- Any matrix  $W \in \mathbb{R}^{m \times n}$  with  $\text{rank}(W) \leq K$  can be expressed as  $W = XY$  where  $X \in \mathbb{R}^{m \times K}$  and  $Y \in \mathbb{R}^{K \times n}$ .

## New model

$$\min_{X,Y,Z} \frac{1}{2} \|XY - Z\|_F^2 \quad \text{s.t.} \quad Z_{ij} = M_{ij}, \forall (i,j) \in \Omega$$

- **Advantage: SVD is no longer needed!**
- Related work: the solver `OptSpace` based on optimization on manifold

# Nonlinear Gauss-Seidel scheme

First variant of alternating minimization:

$$\begin{aligned}X_+ &\leftarrow ZY^\dagger \equiv ZY^\top (YY^\top)^\dagger, \\Y_+ &\leftarrow (X_+)^\dagger Z \equiv (X_+^\top X_+)^\dagger (X_+^\top Z), \\Z_+ &\leftarrow X_+ Y_+ + \mathcal{P}_\Omega(M - X_+ Y_+).\end{aligned}$$

Let  $\mathcal{P}_A$  be the orthogonal projection onto the range space  $\mathcal{R}(A)$

- $X_+ Y_+ = (X_+ (X_+^\top X_+)^{-1} X_+^\top) Z = \mathcal{P}_{X_+} Z$
- One can verify that  $\mathcal{R}(X_+) = \mathcal{R}(ZY^\top)$ .
- $X_+ Y_+ = \mathcal{P}_{ZY^\top} Z = ZY^\top (YZ^\top ZY^\top)^{-1} (YZ^\top) Z$ .
- idea: modify  $X_+$  or  $Y_+$  to obtain the same product  $X_+ Y_+$



# Nonlinear Gauss-Seidel scheme

Second variant of alternating minimization:

$$\begin{aligned}X_+ &\leftarrow ZY^\top, \\Y_+ &\leftarrow (X_+)^{\dagger}Z \equiv (X_+^\top X_+)^{\dagger}(X_+^\top Z), \\Z_+ &\leftarrow X_+Y_+ + \mathcal{P}_\Omega(M - X_+Y_+).\end{aligned}$$

Third variant of alternating minimization:  $V = \text{orth}(ZY^\top)$

$$\begin{aligned}X_+ &\leftarrow V, \\Y_+ &\leftarrow V^\top Z, \\Z_+ &\leftarrow X_+Y_+ + \mathcal{P}_\Omega(M - X_+Y_+).\end{aligned}$$

# Sparse and low-rank matrix separation


- Given a matrix  $M$ , we want to find a low rank matrix  $W$  and a sparse matrix  $E$ , so that  $W + E = M$ .
- Convex approximation:

$$\min_{W,E} \|W\|_* + \mu\|E\|_1, \text{ s.t. } W + E = M$$

- Robust PCA

# Video separation

- Partition the video into moving and static parts



Convex approximation:

$$\min_{W, E} \|W\|_* + \mu\|E\|_1, \text{ s.t. } W + E = M$$

Augmented Lagrangian function:

$$L(W, E, \Lambda) := \|W\|_* + \mu\|E\|_1 + \langle \Lambda, W + E - M \rangle + \frac{1}{2\beta} \|W + E - M\|_F^2$$

Alternating direction Augmented Lagrangian method

$$\begin{aligned} W^{j+1} &:= \arg \min_W L(W, E^j, \Lambda^j), \\ E^{j+1} &:= \arg \min_E L(W^{j+1}, E, \Lambda^j), \\ \Lambda^{j+1} &:= \Lambda^j + \frac{\gamma}{\beta} (W^{j+1} + E^{j+1} - M). \end{aligned}$$

# W-subproblem

Convex approximation:

$$\begin{aligned} W^{j+1} &:= \arg \min_W L(W, E^j, \Lambda^j) \\ &= \arg \min_W \|W\|_* + \frac{1}{2\beta} \|W - (M - E^j - \beta\Lambda^j)\|_F^2 \\ &= S_\beta(M - E^j - \beta\Lambda^j) := U \text{Diag}(s_\beta(\sigma)) V^\top \end{aligned}$$

- SVD:  $M - E^j - \beta\Lambda^j = U \text{Diag}(\sigma) V^\top$
- Thresholding operator:


$$s_\nu(x) := \bar{x}, \text{ with } \bar{x}_i = \begin{cases} x_i - \nu, & \text{if } x_i - \nu > 0 \\ 0, & \text{o.w.} \end{cases}$$

# E-subproblem

Convex approximation:

$$\begin{aligned} W^{j+1} &:= \arg \min_E L(W^{j+1}, E, \Lambda^j) \\ &= \arg \min_E \|E\|_1 + \frac{1}{2\beta\mu} \|E - (M - W^{j+1} - \beta\Lambda^j)\|_F^2 \\ &= s_{\beta\mu}(M - W^{j+1} - \beta\Lambda^j) \end{aligned}$$

$$\begin{aligned} s_\nu(y) : &= \arg \min_{x \in \mathbb{R}} \nu \|x\|_1 + \frac{1}{2} \|x - y\|_2^2 \\ &= \begin{cases} y - \nu \operatorname{sgn}(y), & \text{if } |y| > \nu \\ 0, & \text{otherwise} \end{cases} \end{aligned}$$



# Low-rank factorization model for matrix separation

- Consider the model

$$\min_{Z,S} \|S\|_1 \quad \text{s.t.} \quad Z + S = D, \text{rank}(Z) \leq K$$

- Low-rank factorization:  $Z = UV$

$$\min_{U,V,Z} \|Z - D\|_1 \quad \text{s.t.} \quad UV - Z = 0$$

- Only the entries  $D_{ij}$ ,  $(i,j) \in \Omega$ , are given.  $\mathcal{P}_\Omega(D)$  is the projection of  $D$  onto  $\Omega$ .

## New model

$$\min_{U,V,Z} \|\mathcal{P}_\Omega(Z - D)\|_1 \quad \text{s.t.} \quad UV - Z = 0$$

- Advantage: SVD is no longer needed!

Consider:

$$\min_{U,V,Z} \quad \|\mathcal{P}_\Omega(Z - D)\|_1 \quad \text{s.t.} \quad UV - Z = 0$$

Introduce the augmented Lagrangian function

$$\mathcal{L}_\beta(U, V, Z, \Lambda) = \|\mathcal{P}_\Omega(Z - D)\|_1 + \langle \Lambda, UV - Z \rangle + \frac{\beta}{2} \|UV - Z\|_F^2,$$

Alternating direction augmented Lagrangian framework (Bregman):

$$U^{j+1} := \arg \min_{U \in \mathbb{R}^{m \times k}} \mathcal{L}_\beta(U, V^j, Z^j, \Lambda^j),$$

$$V^{j+1} := \arg \min_{V \in \mathbb{R}^{k \times n}} \mathcal{L}_\beta(U^{j+1}, V, Z^j, \Lambda^j),$$

$$Z^{j+1} := \arg \min_{Z \in \mathbb{R}^{m \times n}} \mathcal{L}_\beta(U^{j+1}, V^{j+1}, Z, \Lambda^j),$$

$$\Lambda^{j+1} := \Lambda^j + \gamma \beta (U^{j+1} V^{j+1} - Z^{j+1}).$$



# ADMM subproblems

- Let  $B = Z - \Lambda/\beta$ , then

$$U_+ = BV^\top(VV^\top)^\dagger \text{ and } V_+ = (U_+^\top U_+)^\dagger U_+^\top B$$

Since  $U_+ V_+ = U_+ (U_+^\top U_+)^\dagger U_+^\top B = \mathcal{P}_{U_+} B$ , then:

$$Q := \text{orth}(BV^\top), \quad U_+ = Q \text{ and } V_+ = Q^\top B$$

- Variable  $Z$ :

$$\mathcal{P}_\Omega(Z_+) = \mathcal{P}_\Omega \left( \mathcal{S} \left( U_+ V_+ - D + \frac{\Lambda}{\beta}, \frac{1}{\beta} \right) + D \right)$$

$$\mathcal{P}_{\Omega^c}(Z_+) = \mathcal{P}_{\Omega^c} \left( U_+ V_+ + \frac{\Lambda}{\beta} \right)$$

# Nonnegative matrix factorization completion

Model problem:

$$\begin{array}{ll} \min & \|\mathcal{P}_\Omega(XY - M)\|_F \\ \text{s.t.} & X \geq 0, Y \geq 0, \end{array} \iff \begin{array}{ll} \min & \frac{1}{2}\|XY - Z\|_F^2 \\ \text{s.t.} & X = U, Y = V, \\ & U \geq 0, V \geq 0, \\ & \mathcal{P}_\Omega(Z - M) = 0 \end{array}$$

Augmented Lagrangian function:

$$\begin{aligned} \mathcal{L}_A(X, Y, Z, U, V, \Lambda, \Pi) = & \frac{1}{2}\|XY - Z\|_F^2 + \Lambda \bullet (X - U) \\ & + \Pi \bullet (Y - V) + \frac{\alpha}{2}\|X - U\|_F^2 + \frac{\beta}{2}\|Y - V\|_F^2, \end{aligned}$$

where  $A \bullet B := \sum_{i,j} a_{ij}b_{ij}$ .

$$X_{k+1} = (Z_k Y_k^T + \alpha U_k - \Lambda_k)(Y_k Y_k^T + \alpha I)^{-1}, \quad (3a)$$

$$Y_{k+1} = (X_{k+1}^T X_{k+1} + \beta I)^{-1}(X_{k+1}^T Z_k + \beta V_k - \Pi_k), \quad (3b)$$

$$Z_{k+1} = X_{k+1} Y_{k+1} + \mathcal{P}_\Omega(M - X_{k+1} Y_{k+1}), \quad (3c)$$

$$U_{k+1} = \mathcal{P}_+(X_{k+1} + \Lambda_k/\alpha), \quad (3d)$$

$$V_{k+1} = \mathcal{P}_+(Y_{k+1} + \Pi_k/\beta), \quad (3e)$$

$$\Lambda_{k+1} = \Lambda_k + \gamma\alpha(X_{k+1} - U_{k+1}), \quad (3f)$$

$$\Pi_{k+1} = \Pi_k + \gamma\beta(Y_{k+1} - V_{k+1}), \quad (3g)$$

where  $\gamma \in (0, 1.618)$  and  $(\mathcal{P}_+(A))_{ij} = \max\{a_{ij}, 0\}$ .

# Sparse covariance selection (A. d'Aspremont)

We estimate a covariance matrix  $\Sigma$  from empirical data

- Infer **independence** relationships between variables
- Given  $m + 1$  observations  $x_i \in \mathbb{R}^n$  on  $n$  random variables, compute  $S := \frac{1}{m} \sum_{i=1}^{m+1} (x_i - \bar{x})(x_i - \bar{x})$
- Choose a symmetric subset  $I$  of matrix coefficients and denote by  $J$  the complement
- Choose a covariance matrix  $\hat{\Sigma}$  such that
  - $\hat{\Sigma}_{ij} = S_{ij}$  for all  $(i, j) \in I$
  - $\hat{\Sigma}_{ij}^{-1} = 0$  for all  $(i, j) \in J$
- Benefits: maximum entropy, maximum likelihood, existence and uniqueness
- Applications: Gene expression data, speech recognition and finance

# Maximum likelihood estimation

Consider estimation:

$$\max_{X \in S^n} \log \det X - \text{Tr}(SX) - \rho \|X\|_0$$

Convex relaxations:

$$\max_{X \in S^n} \log \det X - \text{Tr}(SX) - \rho \|X\|_1,$$

whose dual problem is:

$$\max \log \det W \quad \text{s.t.} \quad \|W - S\|_\infty \leq \lambda$$

Zhaosong Lu (*smooth optimization approach for sparse covariance selection*) consider

$$\begin{aligned} \max \quad & \log \det X - \text{Tr}(SX) - \rho \|X\|_1 \\ \text{s.t.} \quad & \mathcal{X} := \{X \in S^n : \beta I \succeq X \succeq \alpha I\}, \end{aligned}$$

which is equivalent to ( $\mathbf{U} := \{U \in S^n : |U_{ij}| \leq 1, \forall ij\}$ )

$$\max_{X \in \mathcal{X}} \min_{U \in \mathbf{U}} \log \det X - \langle S + \rho U, X \rangle$$

Let  $f(U) := \max_{X \in \mathcal{X}} \log \det X - \langle S + \rho U, X \rangle$

- $\log \det X$  is strongly concave on  $\mathcal{X}$
- $f(U)$  is continuous differentiable
- $\nabla f(U)$  is Lipschitz cont. with  $L = \rho\beta^2$

Therefore, APG can be applied to the dual problem

$$\min_{U \in \mathbf{U}} f(U)$$

# Extension

Consider

$$\max_{x \in \mathcal{X}} g(x) := \min_{u \in \mathbf{U}} \phi(x, u)$$

Assume:

- $\phi(x, u)$  is a cont. fun. which is strictly concave in  $x \in \mathcal{X}$  for every fixed  $u \in \mathbf{U}$ , and convex diff. in  $u \in \mathbf{U}$  for every fixed  $x \in \mathcal{X}$ . Then  $f(u) := \max_{x \in \mathcal{X}} \phi(x, u)$  is diff.
- $\nabla f(u)$  is Lipschitz cont.

Then

- the primal and the dual  $\min_{u \in U} f(u)$  are both solvable and have the same optimal value;
- Nesterov's smooth minimization approach can be applied to the dual

# Nesterov's smoothing technique

Consider

$$\max_{x \in \mathcal{X}} \min_{u \in \mathbf{U}} \phi(x, u)$$

Question: What if the assumptions do not hold?

- Add a strictly convex function  $\mu d(u)$  to the obj. fun.

$$g(u) := \arg \min_{u \in \mathbf{U}} \phi(x, u) + \mu d(u)$$

- $g(u)$  is differentiable
- Apply Nesterov's smooth minimization
- Complexity of finding a  $\epsilon$ -suboptimal point:  $O(\frac{1}{\epsilon})$  iterations
- Other smooth technique?